arXiv:1604.02950v1 [math.RA] 11 Apr 2016

Rota-Baxter coalgebras and Rota-Baxter bialgebras

Tianshui Ma* **and Linlin Liu**

[1]Department of Mathematics, School of Mathematics and Information Science,
Henan Normal University, Xinxiang 453007, China

[2]Henan Engineering Laboratory for Big Data Statistical Analysis and Optimal Control,
School of Mathematics and Information Sciences, Henan Normal University,
Xinxiang 453007, China

E-mail: matianshui@yahoo.com (T. Ma); liulinlin2016@163.com (L. Liu)

ABSTRACT

We give a construction of Rota-Baxter coalgebras from Hopf module coalgebras and also derive the structures of the pre-Lie coalgebras via Rota-Baxter coalgebras of different weight. Finally, the notion of Rota-Baxter bialgebra is introduced and some examples are provided.

Key words: Rota-Baxter bialgebras, Radford biproduct, Rota-Baxter coalgebras.

2010 Mathematics Subject Classification: 16T05, 16W99

1 Introduction

A Rota-Baxter algebra, also called a Baxter algebra, is an associative algebra with a linear operator which generalizes the algebra of continuous functions with the integral operator. More precisely, for a given field K and $\lambda \in K$, a Rota-Baxter K-algebra (of weight λ) is a K-algebra R together with a K-linear map $P : R \longrightarrow R$ such that

$$P(x)P(y) = P(xP(y)) + P(P(x)y) + \lambda P(xy) \tag{1}$$

for all $x, y \in R$. Such a linear operator is called a Rota-Baxter operator (of weight λ). Rota-Baxter algebras were introduced in [21] in the context of differential operators on commutative Banach algebras and since [4] intensively studied in probability and combinatorics, and more recently in mathematical physics, such as dendriform algebras

*Corresponding author

(see [3, 6, 7, 11]), free Rota-Baxter algebras (see [5, 8, 9, 13, 14]), Lie algebras (see [2, 18]), multiple zeta values (see [8, 15]) and Connes-Kreimer renormalization theory in quantum field theory (see [10]), etc. One can refer to the book [12] for the detailed theory of Rota-Baxter algebras.

In 2014, Jian used Yetter-Drinfeld module algebras to construct Hopf module algebras, and then the corresponding Rota-Baxter algebras were derived (see [16]). Based on the dual method in the Hopf algebra theory, Jian and Zhang in [17] defined the notion of Rota-Baxter coalgebras and also provided various examples of the new object, including constructions by group-like elements and by smash coproduct.

In this paper, we will continue to investigate the properties of Rota-Baxter coalgebras. We give a construction of Rota-Baxter coalgebras from Hopf module coalgebras and also derive the structures of the pre-Lie coalgebras via Rota-Baxter coalgebras of different weight. Finally, the notion of Rota-Baxter bialgebra is introduced and some examples are provided.

2 Preliminaries

Throughout this paper, we assume that all vector spaces, algebras, coalgebras and tensor products are defined over a field K. An algebra is always assumed to be associative, but not necessarily unital. A coalgebra is always assumed to be coassociative, but not necessarily counital. Now, let C be a coalgebra. We use Sweedler's notation for the comultiplication (see [19]): $\Delta(c) = c_1 \otimes c_2$ for any $c \in C$. Denote the category of left H-comodules by ${}^H\mathrm{Mod}$. For $(M, \rho_L) \in {}^H\mathrm{Mod}$, write: $\rho_L(x) = x_{(-1)} \otimes x_{(0)} \in H \otimes M$, for all $x \in M$. Denote the category of right H-comodules by Mod^H. For $(M, \rho_R) \in \mathrm{Mod}^H$, write: $\rho_R(x) = x_{(0)} \otimes x_{(1)} \in M \otimes H$, for all $x \in M$. Given a K-space M, we write id_M for the identity map on M.

In what follows, we recall some useful definitions and results which will be used later (see [12, 16, 17, 19]).

Hopf module (see [19]): A right (resp. left) H-Hopf module is a vector space M equipped simultaneously with a right (resp. left) H-module structure and a right (resp. left) H-comodule structure ρ_R (resp. ρ_L) such that

$$\rho_R(m \cdot h) = \rho_R(m)\Delta(h). \tag{2}$$

(resp. $\rho_L(h \cdot m) = \Delta(h)\rho_L(m)$) for all $h \in H$ and $m \in M$.

Remark Let M be a right H-Hopf module. We denote by M^R the subspace of right coinvariants, i.e.,

$$M^R = \{m \in M \mid \rho_R(m) = m \otimes 1_H\}.$$

Then the map $P_R : M \longrightarrow M$ given by $P_R(m) = m_{(0)} \cdot S(m_{(1)})$ for $m \in M$ is the projection from M onto M^R. Similarly, for a left H-Hopf module, we can define the subspace M^L of left coinvariants, i.e.,

$$M^L = \{m \in M \mid \rho_L(m) = 1_H \otimes m\},$$

and have the projection formula

$$P_L(m) = S(m_{(-1)}) \cdot m_{(0)}.$$

Hopf module algebra (see [16]): A right H-Hopf module algebra is a right H-Hopf module M together with an associative multiplication $\mu : M \otimes M \longrightarrow M$ (write $\mu(m \otimes m') = mm'$) such that for any $h \in H$ and $m, m' \in M$,

$$(mm') \cdot h = m(m' \cdot h), \tag{3}$$

$$\rho_R(mm') = m_{(0)}m'_{(0)} \otimes m_{(1)}m'_{(1)}. \tag{4}$$

Remarks (1) The compatibility conditions between μ and the Hopf module structure mean that μ is both a module morphism and a comodule morphism,where the right H-module action and right H-comodule coaction on $M \otimes M$ are given by

$$M \otimes M \otimes H \longrightarrow M \otimes M \quad (m \otimes m' \otimes h \mapsto m \otimes m' \cdot h),$$

$$M \otimes M \longrightarrow M \otimes M \otimes H \quad (m \otimes m' \mapsto m_{(0)} \otimes m'_{(0)} \otimes m_{(1)}m'_{(1)})$$

for all $h \in H$ and $m, m' \in M$.

(2) We can define a left H-Hopf module algebra by satisfying the compatibility conditions $h \cdot (mm') = (h \cdot m)m'$ and $\rho_L(mm') = m_{(-1)}m'_{(-1)} \otimes m_{(0)}m'_{(0)}$.

Rota-Baxter coalgebra (see [17]): Let γ be an element in K. A pair (C, Q) is called a Rota-Baxter coalgebra of weight γ if C is a coalgebra and Q is a linear endomorphism of C satisfying that

$$(Q \otimes Q)\Delta = (id \otimes Q)\Delta Q + (Q \otimes id)\Delta Q + \gamma \Delta Q,$$

that is, for all $c \in C$,

$$Q(c_1) \otimes Q(c_2) = Q(c)_1 \otimes Q(Q(c)_2) + Q(Q(c)_1) \otimes Q(c)_2 + \gamma Q(c)_1 \otimes Q(c)_2. \tag{5}$$

The map Q is called a Rota-Baxter operator.

A Rota-Baxter coalgebra is called idempotent if the Rota-Baxter operator Q satisfies $Q^2 = Q$.

Yetter-Drinfeld category (see [19]): A left H-Yetter-Drinfeld module is a vector space V equipped with a left H-module structure $\cdot$ and a left H-comodule structure ρ_L such that for all $h \in H$ and $v \in V$,

$$h_1 v_{(-1)} \otimes h_2 \cdot v_{(0)} = (h_1 \cdot v)_{(-1)} h_2 \otimes (h_1 \cdot v)_{(0)}. \tag{6}$$

The category of left H-Yetter-Drinfeld modules, denoted by ${}^H_H\mathcal{YD}$, consists of the following data: Its objects are left H-Yetter-Drinfeld modules and morphisms are linear maps which are both module and comodule morphisms.

Remarks (1) A coalgebra (C, Δ) in ${}^H_H\mathcal{YD}$ is a coassociative coalgebra such that the underlying space C is an object in ${}^H_H\mathcal{YD}$ and (C, Δ) is both a module-coalgebra and a comodule-coalgebra. More precisely, for all $h \in H$ and $c \in C$ we have

$$\Delta(h \cdot c) = h_1 \cdot c_1 \otimes h_2 \cdot c_2, \tag{7}$$

$$c_{(-1)} \otimes c_{(0)1} \otimes c_{(0)2} = c_{1(-1)} c_{2(-1)} \otimes c_{1(0)} \otimes c_{2(0)}. \tag{8}$$

(2) Let (C, Δ) be a coalgebra in ${}^H_H\mathcal{YD}$. The smash coproduct coalgebra $C \times H$ is defined to be $C \otimes H$ as a vector space and

$$\Delta(c \otimes h) = c_1 \otimes c_{2(-1)} h_1 \otimes c_{2(0)} \otimes h_2. \tag{9}$$

3 Construction of Rota-Baxter coalgebras

In this section, we give a construction of Rota-Baxter coalgebras via Hopf module coalgebras.

Definition 3.1 A right H-Hopf module coalgebra is a right H-Hopf module M together with coassociative comultiplication $\Delta : M \longrightarrow M \otimes M$ $(\Delta(m) = m_1 \otimes m_2)$ such that for any $h \in H$ and $m \in M$,

$$m_{(0)1} \otimes m_{(0)2} \otimes m_{(1)} = m_1 \otimes m_{2(0)} \otimes m_{2(1)}, \tag{10}$$

$$\Delta(m \cdot h) = m_1 \cdot h_1 \otimes m_2 \cdot h_2. \tag{11}$$

Remarks (1) The compatibility conditions between Δ and the Hopf module structure mean that Δ is both a module morphism and a comodule morphism, where the right H-module action and right H-comodule coaction on $M \otimes M$ are given by

$$M \otimes M \otimes H \longrightarrow M \otimes M \quad (m \otimes m' \otimes h \mapsto m \cdot h_1 \otimes m' \cdot h_2),$$

$$M \otimes M \longrightarrow M \otimes M \otimes H \quad (m \otimes m' \mapsto m \otimes m'_{(0)} \otimes m'_{(1)})$$

for all $h \in H$ and $m, m' \in M$.

(2) We can define a left H-Hopf module coalgebra by satisfying the compatibility conditions:

$$m_{(-1)} \otimes m_{(0)1} \otimes m_{(0)2} = m_{1(-1)} \otimes m_{1(0)} \otimes m_2,$$

and

$$\Delta(h \cdot m) = h_1 \cdot m_1 \otimes h_2 \cdot m_2,$$

where $h \in H$ and $m \in M$.

Theorem 3.2 Let (C, Δ) be a right H-Hopf module coalgebra. Then (C, P_R) is a Rota-Baxter coalgebra of weight -1.

Proof. Since C is a right H-comodule, for all $c \in C$ we have

$$c_{(0)} \otimes c_{(1)1} \otimes c_{(1)2} = c_{(0)(0)} \otimes c_{(0)(1)} \otimes c_{(1)}. \tag{12}$$

We check the condition (5) for P_R as follows:

$$\begin{aligned}
& ((id \otimes P_R)\Delta P_R + (P_R \otimes id)\Delta P_R - \Delta P_R)(c) \\
= \; & (id \otimes P_R)\Delta P_R(c) + (P_R \otimes id)\Delta P_R(c) - \Delta P_R(c) \\
= \; & (id \otimes P_R)\Delta(c_{(0)} \cdot S(c_{(1)})) + (P_R \otimes id)\Delta(c_{(0)} \cdot S(c_{(1)})) - \Delta(c_{(0)} \cdot S(c_{(1)})) \\
= \; & (c_{(0)} \cdot S(c_{(1)}))_1 \otimes P_R((c_{(0)} \cdot S(c_{(1)}))_2) + P_R((c_{(0)} \cdot S(c_{(1)}))_1) \otimes (c_{(0)} \cdot S(c_{(1)}))_2 \\
& -(c_{(0)} \cdot S(c_{(1)}))_1 \otimes (c_{(0)} \cdot S(c_{(1)}))_2 \\
\overset{(11)}{=} \; & c_{(0)1} \cdot S(c_{(1)2}) \otimes (c_{(0)2} \cdot S(c_{(1)1}))_{(0)} \cdot S((c_{(0)2} \cdot S(c_{(1)1}))_{(1)}) + (c_{(0)1} \cdot S(c_{(1)2}))_{(0)} \\
& \cdot S((c_{(0)1} \cdot S(c_{(1)2}))_{(1)}) \otimes c_{(0)2} \cdot S(c_{(1)1}) - c_{(0)1} \cdot S(c_{(1)2}) \otimes c_{(0)2} \cdot S(c_{(1)1}) \\
\overset{(2)}{=} \; & c_{(0)1} \cdot S(c_{(1)2}) \otimes (c_{(0)2(0)} \cdot S(c_{(1)1})_1) \cdot S(c_{(0)2(1)} S(c_{(1)1})_2) + c_{(0)1(0)} \cdot S(c_{(1)2})_1 \\
& \cdot S(c_{(0)1(1)} S(c_{(1)2})_2) \otimes c_{(0)2} \cdot S(c_{(1)1}) - c_{(0)1} \cdot S(c_{(1)2}) \otimes c_{(0)2} \cdot S(c_{(1)1}) \\
= \; & c_{(0)1} \cdot S(c_{(1)2}) \otimes c_{(0)2(0)} \cdot ((S(c_{(1)1})_1 S(S(c_{(1)1})_2)) S(c_{(0)2(1)})) + c_{(0)1(0)} \cdot ((S(c_{(1)2})_1 \\
& S(S(c_{(1)2})_2)) S(c_{(0)1(1)})) \otimes c_{(0)2} \cdot S(c_{(1)1}) - c_{(0)1} \cdot S(c_{(1)2}) \otimes c_{(0)2} \cdot S(c_{(1)1}) \\
= \; & c_{(0)1} \cdot S(c_{(1)2}) \otimes c_{(0)2(0)} \cdot \varepsilon(S(c_{(1)1})) S(c_{(0)2(1)}) + c_{(0)1(0)} \cdot S(c_{(0)1(1)}) \\
& \otimes c_{(0)2} \cdot S(c_{(1)1}) \varepsilon(S(c_{(1)2})) - c_{(0)1} \cdot S(c_{(1)2}) \otimes c_{(0)2} \cdot S(c_{(1)1}) \\
= \; & c_{(0)1} \cdot S(c_{(1)}) \otimes c_{(0)2(0)} \cdot S(c_{(0)2(1)}) + c_{(0)1(0)} \cdot S(c_{(0)1(1)}) \otimes c_{(0)2} \cdot S(c_{(1)}) \\
& -c_{(0)1} \cdot S(c_{(1)2}) \otimes c_{(0)2} \cdot S(c_{(1)1}) \\
\overset{(10)}{=} \; & c_{(0)(0)1} \cdot S(c_{(1)}) \otimes c_{(0)(0)2} \cdot S(c_{(0)(1)}) + c_{(0)1(0)} \cdot S(c_{(0)1(1)}) \otimes c_{(0)2} \cdot S(c_{(1)}) \\
& -c_{(0)1} \cdot S(c_{(1)2}) \otimes c_{(0)2} \cdot S(c_{(1)1})
\end{aligned}$$

$$
\begin{aligned}
&\overset{(12)}{=} && c_{(0)1}\cdot S(c_{(1)2})\otimes c_{(0)2}\cdot S(c_{(1)1})+c_{(0)1(0)}\cdot S(c_{(0)1(1)})\otimes c_{(0)2}\cdot S(c_{(1)})\\
&&& -c_{(0)1}\cdot S(c_{(1)2})\otimes c_{(0)2}\cdot S(c_{(1)1})\\
&= && c_{(0)1(0)}\cdot S(c_{(0)1(1)})\otimes c_{(0)2}\cdot S(c_{(1)})\\
&\overset{(10)}{=} && c_{1(0)}\cdot S(c_{1(1)})\otimes c_{2(0)}\cdot S(c_{2(1)})=(P_R\otimes P_R)\Delta(c),
\end{aligned}
$$

finishing the proof. □

Remark Let (C,Δ) be a left H-Hopf module coalgebra. Then (C,P_L) is a Rota-Baxter coalgebra of weight -1.

Example 3.3 Let C be a bialgebra and H a Hopf algebra with the antipode S. Suppose that there are two bialgebra maps: $i:H\longrightarrow C$ and $\pi:C\longrightarrow H$ such that $\pi\circ i=id_H$, i.e., C is a bialgebra with a projection (see [20]). Set $\Pi=id_C\star(i\circ S\circ\pi)$, where $\star$ is the convolution product on $\mathrm{End}(C)$. The right H-module structure and right H-comodule structure on C are given by the following:

$$c\cdot h=ci(h),$$

$$\rho_R(c)=c_1\otimes\pi(c_2),$$

for all $c\in C$ and $h\in H$. Then (C,Π) is a Rota-Baxter coalgebra of weight -1.

Proof. For all $h\in H$ and $c\in C$, firstly C is a right H-Hopf module with the above module structure and comodule structure, since

$$
\begin{aligned}
\rho_R(c\cdot h) &= \rho_R(ci(h))=(ci(h))_1\otimes\pi((ci(h))_2)\\
&= c_1i(h_1)\otimes\pi(c_2)h_2=c_1\cdot h_1\otimes\pi(c_2)h_2=\rho_R(c)\Delta(h).
\end{aligned}
$$

Secondly, we have

$$
\begin{aligned}
(\Delta\otimes id)\rho_R(c) &= (\Delta\otimes id)(c_1\otimes\pi(c_2))=c_{11}\otimes c_{12}\otimes\pi(c_2)\\
&= c_1\otimes c_{21}\otimes\pi(c_{22})=(id\otimes\rho_R)\Delta(c)
\end{aligned}
$$

and $\Delta(c\cdot h)=c_1i(h_1)\otimes c_2i(h_2)=c_1\cdot h_1\otimes c_2\cdot h_2$. Thus (C,Δ) is a right H-Hopf module coalgebra.

Finally, by [20, Theorem 3], Π is the projection from C to C^R. Therefore, we get the conclusion by Theorem 3.2. □

Next we investigate the Hopf module coalgebra structure on the smash coproduct coalgebra.

Theorem 3.4 Let (C,Δ) be a coalgebra in ${}^H_H\mathcal{YD}$. Then: (1) The smash coproduct coalgebra $C\times H$ is a right H-Hopf module coalgebra with the following right H-Hopf module structure: for all $h,x\in H$ and $c\in C$,

$$(c\otimes h)\cdot x=c\otimes hx,$$

$$\rho_R(c \otimes h) = (c \otimes h_1) \otimes h_2.$$

Hence the map P_R given by

$$P_R(c \otimes h) = c \otimes \varepsilon(h)1_H$$

is an idempotent Rota-Baxter operator of weight -1.

(2) The smash coproduct coalgebra $C \times H$ is a left H-Hopf module coalgebra with the following left H-Hopf module structure:

$$x \cdot (c \otimes h) = x_1 \cdot c \otimes x_2 h,$$

$$\rho_L(c \otimes h) = c_{(-1)}h_1 \otimes (c_{(0)} \otimes h_2)$$

Hence the map P_L given by

$$P_L(c \otimes h) = S(c_{(-1)2}h_2) \cdot c_{(0)} \otimes S(c_{(-1)1}h_1)h_3$$

is an idempotent Rota-Baxter operator of weight -1.

Proof. In what follows, we only prove the compatibility conditions, and the constructions of the right and left H-Hopf module structures given above are obvious.

(1) For all $c \in C$ and $h, x \in H$, we have

$$\begin{aligned}
\Delta((c \otimes h) \cdot x) &= \Delta(c \otimes hx) \overset{(9)}{=} c_1 \otimes c_{2(-1)}(hx)_1 \otimes c_{2(0)} \otimes (hx)_2 \\
&= c_1 \otimes (c_{2(-1)}h_1)x_1 \otimes c_{2(0)} \otimes h_2 x_2 \\
&= (c_1 \otimes c_{2(-1)}h_1) \cdot x_1 \otimes (c_{2(0)} \otimes h_2) \cdot x_2 \\
&= (c \otimes h)_1 \cdot x_1 \otimes (c \otimes h)_2 \cdot x_2.
\end{aligned}$$

and

$$\begin{aligned}
(id \otimes \rho_R)\Delta(c \otimes h) &\overset{(9)}{=} (id \otimes \rho_R)(c_1 \otimes c_{2(-1)}h_1 \otimes c_{2(0)} \otimes h_2) \\
&= c_1 \otimes c_{2(-1)}h_1 \otimes c_{2(0)} \otimes h_{21} \otimes h_{22} \\
&= c_1 \otimes c_{2(-1)}h_{11} \otimes c_{2(0)} \otimes h_{12} \otimes h_2 \\
&= (\Delta \otimes id)(c \otimes h_1 \otimes h_2) = (\Delta \otimes id)\rho_R(c \otimes h).
\end{aligned}$$

(2) Since C is a left H-comodule, for all $c \in C$ we have

$$c_{(-1)} \otimes c_{(0)(-1)} \otimes c_{(0)(0)} = c_{(-1)1} \otimes c_{(-1)2} \otimes c_{(0)}. \tag{13}$$

While

$$\begin{aligned}
\Delta(x\cdot(c\otimes h)) &= \Delta(x_1\cdot c\otimes x_2h)\\
&\overset{(9)}{=} (x_1\cdot c)_1\otimes(x_1\cdot c)_{2(-1)}(x_2h)_1\otimes(x_1\cdot c)_{2(0)}\otimes(x_2h)_2\\
&\overset{(7)}{=} x_{11}\cdot c_1\otimes(x_{12}\cdot c_2)_{(-1)}(x_{21}h_1)\otimes(x_{12}\cdot c_2)_{(0)}\otimes x_{22}h_2\\
&= x_1\cdot c_1\otimes((x_2\cdot c_2)_{(-1)}x_3)h_1\otimes(x_2\cdot c_2)_{(0)}\otimes x_4h_2\\
&\overset{(6)}{=} x_1\cdot c_1\otimes(x_2c_{2(-1)})h_1\otimes x_3\cdot c_{2(0)}\otimes x_4h_2\\
&= x_{11}\cdot c_1\otimes x_{12}(c_{2(-1)}h_1)\otimes x_{21}\cdot c_{2(0)}\otimes x_{22}h_2\\
&= x_1\cdot(c_1\otimes c_{2(-1)}h_1)\otimes x_2\cdot(c_{2(0)}\otimes h_2)\\
&= x_1\cdot(c\otimes h)_1\otimes x_2\cdot(c\otimes h)_2,
\end{aligned}$$

and

$$\begin{aligned}
(id\otimes\Delta)\rho_L(c\otimes h) &= (c\otimes h)_{(-1)}\otimes(c\otimes h)_{(0)1}\otimes(c\otimes h)_{(0)2}\\
&= c_{(-1)}h_1\otimes(c_{(0)}\otimes h_2)_1\otimes(c_{(0)}\otimes h_2)_2\\
&= c_{(-1)}h_1\otimes c_{(0)1}\otimes c_{(0)2(-1)}h_{21}\otimes c_{(0)2(0)}\otimes h_{22}\\
&= c_{(-1)}h_1\otimes c_{(0)1}\otimes c_{(0)2(-1)}h_2\otimes c_{(0)2(0)}\otimes h_3\\
&\overset{(8)}{=} (c_{1(-1)}c_{2(-1)})h_1\otimes c_{1(0)}\otimes c_{2(0)(-1)}h_2\otimes c_{2(0)(0)}\otimes h_3\\
&\overset{(13)}{=} (c_{1(-1)}c_{2(-1)1})h_1\otimes c_{1(0)}\otimes c_{2(-1)2}h_2\otimes c_{2(0)}\otimes h_3\\
&= c_{1(-1)}(c_{2(-1)}h_1)_1\otimes c_{1(0)}\otimes(c_{2(-1)}h_1)_2\otimes c_{2(0)}\otimes h_2\\
&= (c_1\otimes c_{2(-1)}h_1)_{(-1)}\otimes(c_1\otimes c_{2(-1)}h_1)_{(0)}\otimes c_{2(0)}\otimes h_2\\
&\overset{(9)}{=} (c\otimes h)_{1(-1)}\otimes(c\otimes h)_{1(0)}\otimes(c\otimes h)_2\\
&= (\rho_L\otimes id)\Delta(c\otimes h),
\end{aligned}$$

thus we complete the proof. □

Example 3.5 The coalgebra H is a coalgebra in ${}^H_H\mathcal{YD}$ with the following left H-Yetter-Drinfeld module structure:

$$x\cdot h = xh,$$

$$\rho(h) = h_1S(h_3)\otimes h_2.$$

Then the smash coproduct on $H\otimes H$ is given by

$$\Delta(h\otimes h') = h_1\otimes(h_2S(h_4))h'_1\otimes h_3\otimes h'_2,$$

the corresponding Rota-Baxter operators P_R and P_L are given by

$$P_R(h\otimes h') = h\otimes 1_H\varepsilon(h')$$

and

$$\begin{aligned} P_L(h \otimes h') &= S((h_1 S(h_3))_2 h'_2) \cdot h_2 \otimes S((h_1 S(h_3))_1 h'_1) h'_3 \\ &= S((h_{12} S(h_{31})) h'_2) \cdot h_2 \otimes S((h_{11} S(h_{32})) h'_1) h'_3 \\ &= S((h_2 S(h_4)) h'_2) h_2 \otimes S((h_1 S(h_5)) h'_1) h'_3. \end{aligned}$$

We recall from [19] that a coquasitriangular (or braided) Hopf algebra is a pair (H, σ), where H is a Hopf algebra and $\sigma \in (H \otimes H)^*$ such that

$$\begin{aligned} &(BR1) \quad \sigma(1, h) = \sigma(h, 1) = \varepsilon(h), \\ &(BR2) \quad \sigma(hh', h'') = \sigma(h, h''_1)\sigma(h', h''_2), \\ &(BR3) \quad \sigma(h, h'h'') = \sigma(h_1, h'')\sigma(h_2, h'), \\ &(BR4) \quad h'_1 h_1 \sigma(h_2, h'_2) = \sigma(h_1, h'_1) h_2 h'_2 \end{aligned}$$

for any $h, h', h'' \in H$.

Example 3.6 Let (H, σ) be a coquasitriangular Hopf algebra. For any H-comodule (M, ρ), we define $\cdot : H \otimes M \longrightarrow M$ by $h \cdot m = \sigma(m_{(-1)}, h) m_{(0)}$. Then $(M, \cdot, \rho)$ is a left H-Yetter-Drinfeld module. By [19], we know that any comodule-coalgebra C over H is moreover a coalgebra in ${}^H_H\mathcal{YD}$. Then Rota-baxter operators P_R and P_L on $C \times H$ are given by

$$P_R(c \otimes h) = c \otimes \varepsilon(h) 1_H$$

and

$$\begin{aligned} P_L(c \otimes h) &= S(c_{(-1)2} h_2) \cdot c_{(0)} \otimes S(c_{(-1)1} h_1) h_3 \\ &= \sigma(c_{(0)(-1)}, S(c_{(-1)2} h_2)) c_{(0)(0)} \otimes S(c_{(-1)1} h_1) h_3 \\ &\overset{(13)}{=} \sigma(c_{(-1)2}, S(c_{(-1)12} h_2)) c_{(0)} \otimes S(c_{(-1)11} h_1) h_3 \\ &= \sigma(c_{(-1)3}, S(c_{(-1)2} h_2)) c_{(0)} \otimes S(c_{(-1)1} h_1) h_3 \\ &= \sigma(c_{(-1)3}, S(h_2) S(c_{(-1)2})) c_{(0)} \otimes S(c_{(-1)1} h_1) h_3 \\ &\overset{(BR3)}{=} \sigma(c_{(-1)3}, S(c_{(-1)2})) \sigma(c_{(-1)4}, S(h_2)) c_{(0)} \otimes S(c_{(-1)1} h_1) h_3 \end{aligned}$$

respectively.

4 From Rota-Baxter coalgebra to pre-Lie coalgebra

In this section, we construct pre-Lie coalgebras from Rota-Baxter coalgebras. First, we give the definition of pre-Lie coalgebra which is dual to the definition of pre-Lie algebra in [1].

Definition 4.1 A pre-Lie coalgebra is (C, Δ) consisting of a linear space C, a linear map $\Delta : C \longrightarrow C \otimes C$ and satisfying

$$\Delta_C - \Phi_{(12)}\Delta_C = 0, \tag{14}$$

where $\Delta_C = (\Delta \otimes id)\Delta - (id \otimes \Delta)\Delta$ and $\Phi_{(12)}(c_1 \otimes c_2 \otimes c_3) = c_2 \otimes c_1 \otimes c_3$.

Theorem 4.2 Let (C, Δ, Q) be a Rota-Baxter coalgebra of weight -1. Define the operation $\widetilde{\Delta}$ on C by

$$\widetilde{\Delta}(c) = Q(c_1) \otimes c_2 - Q(c_2) \otimes c_1 - c_1 \otimes c_2.$$

Then $\widetilde{C} = (C, \widetilde{\Delta})$ is a pre-Lie coalgebra.

Proof. For all $c \in C$, we have

$$\begin{aligned}
\widetilde{\Delta}_C(c) &= (\widetilde{\Delta} \otimes id)\widetilde{\Delta}(c) - (id \otimes \widetilde{\Delta})\widetilde{\Delta}(c) \\
&= (\widetilde{\Delta} \otimes id)(Q(c_1) \otimes c_2 - Q(c_2) \otimes c_1 - c_1 \otimes c_2) \\
&\quad - (id \otimes \widetilde{\Delta})(Q(c_1) \otimes c_2 - Q(c_2) \otimes c_1 - c_1 \otimes c_2) \\
&= \widetilde{\Delta}(Q(c_1)) \otimes c_2 - \widetilde{\Delta}(Q(c_2)) \otimes c_1 - \widetilde{\Delta}(c_1) \otimes c_2 - Q(c_1) \otimes \widetilde{\Delta}(c_2) \\
&\quad + Q(c_2) \otimes \widetilde{\Delta}(c_1) + c_1 \otimes \widetilde{\Delta}(c_2) \\
&= Q(Q(c_1)_1) \otimes Q(c_1)_2 \otimes c_2 - Q(Q(c_1)_2) \otimes Q(c_1)_1 \otimes c_2 - Q(c_1)_1 \otimes Q(c_1)_2 \otimes c_2 \\
&\quad - Q(Q(c_2)_1) \otimes Q(c_2)_2 \otimes c_1 + Q(Q(c_2)_2) \otimes Q(c_2)_1 \otimes c_1 + Q(c_2)_1 \otimes Q(c_2)_2 \otimes c_1 \\
&\quad - Q(c_{11}) \otimes c_{12} \otimes c_2 + Q(c_{12}) \otimes c_{11} \otimes c_2 + c_{11} \otimes c_{12} \otimes c_2 - Q(c_1) \otimes Q(c_{21}) \otimes c_{22} \\
&\quad + Q(c_1) \otimes Q(c_{22}) \otimes c_{21} + Q(c_1) \otimes c_{21} \otimes c_{22} + Q(c_2) \otimes Q(c_{11}) \otimes c_{12} \\
&\quad - Q(c_2) \otimes Q(c_{12}) \otimes c_{11} - Q(c_2) \otimes c_{11} \otimes c_{12} + c_1 \otimes Q(c_{21}) \otimes c_{22} \\
&\quad - c_1 \otimes Q(c_{22}) \otimes c_{21} - c_1 \otimes c_{21} \otimes c_{22} \\
&= Q(Q(c_1)_1) \otimes Q(c_1)_2 \otimes c_2 - Q(Q(c_1)_2) \otimes Q(c_1)_1 \otimes c_2 - Q(c_1)_1 \otimes Q(c_1)_2 \otimes c_2 \\
&\quad - Q(Q(c_2)_1) \otimes Q(c_2)_2 \otimes c_1 + Q(Q(c_2)_2) \otimes Q(c_2)_1 \otimes c_1 + Q(c_2)_1 \otimes Q(c_2)_2 \otimes c_1 \\
&\quad - Q(c_1) \otimes c_2 \otimes c_3 + Q(c_2) \otimes c_1 \otimes c_3 + c_1 \otimes c_2 \otimes c_3 - Q(c_1) \otimes Q(c_2) \otimes c_3 \\
&\quad + Q(c_1) \otimes Q(c_3) \otimes c_2 + Q(c_1) \otimes c_2 \otimes c_3 + Q(c_3) \otimes Q(c_1) \otimes c_2 \\
&\quad - Q(c_3) \otimes Q(c_2) \otimes c_1 - Q(c_3) \otimes c_1 \otimes c_2 + c_1 \otimes Q(c_2) \otimes c_3 \\
&\quad - c_1 \otimes Q(c_3) \otimes c_2 - c_1 \otimes c_2 \otimes c_3 \\
&= Q(Q(c_1)_1) \otimes Q(c_1)_2 \otimes c_2 - Q(Q(c_1)_2) \otimes Q(c_1)_1 \otimes c_2 - Q(c_1)_1 \otimes Q(c_1)_2 \otimes c_2 \\
&\quad - Q(Q(c_2)_1) \otimes Q(c_2)_2 \otimes c_1 + Q(Q(c_2)_2) \otimes Q(c_2)_1 \otimes c_1 + Q(c_2)_1 \otimes Q(c_2)_2 \otimes c_1 \\
&\quad + Q(c_2) \otimes c_1 \otimes c_3 - Q(c_1) \otimes Q(c_2) \otimes c_3 + Q(c_1) \otimes Q(c_3) \otimes c_2 \\
&\quad + Q(c_3) \otimes Q(c_1) \otimes c_2 - Q(c_3) \otimes Q(c_2) \otimes c_1 - Q(c_3) \otimes c_1 \otimes c_2 + c_1 \otimes Q(c_2) \otimes c_3 \\
&\quad - c_1 \otimes Q(c_3) \otimes c_2.
\end{aligned}$$

Then we can verify that the condition (14) holds by using the above equality as follows.

$$
\begin{aligned}
&(\widetilde{\Delta}_C - \Phi_{(12)}\widetilde{\Delta}_C)(c)\\
&\quad = Q(Q(c_1)_1)\otimes Q(c_1)_2\otimes c_2 - Q(Q(c_1)_2)\otimes Q(c_1)_1\otimes c_2 - Q(c_1)_1\otimes Q(c_1)_2\otimes c_2\\
&\qquad - Q(Q(c_2)_1)\otimes Q(c_2)_2\otimes c_1 + Q(Q(c_2)_2)\otimes Q(c_2)_1\otimes c_1 + Q(c_2)_1\otimes Q(c_2)_2\otimes c_1\\
&\qquad + Q(c_2)\otimes c_1\otimes c_3 - Q(c_1)\otimes Q(c_2)\otimes c_3 + Q(c_1)\otimes Q(c_3)\otimes c_2\\
&\qquad + Q(c_3)\otimes Q(c_1)\otimes c_2 - Q(c_3)\otimes Q(c_2)\otimes c_1 - Q(c_3)\otimes c_1\otimes c_2 + c_1\otimes Q(c_2)\otimes c_3\\
&\qquad - c_1\otimes Q(c_3)\otimes c_2 - Q(c_1)_2\otimes Q(Q(c_1)_1)\otimes c_2 + Q(c_1)_1\otimes Q(Q(c_1)_2)\otimes c_2\\
&\qquad + Q(c_1)_2\otimes Q(c_1)_1\otimes c_2 + Q(c_2)_2\otimes Q(Q(c_2)_1)\otimes c_1 - Q(c_2)_1\otimes Q(Q(c_2)_2)\otimes c_1\\
&\qquad - Q(c_2)_2\otimes Q(c_2)_1\otimes c_1 - c_1\otimes Q(c_2)\otimes c_3 + Q(c_2)\otimes Q(c_1)\otimes c_3\\
&\qquad - Q(c_3)\otimes Q(c_1)\otimes c_2 - Q(c_1)\otimes Q(c_3)\otimes c_2 + Q(c_2)\otimes Q(c_3)\otimes c_1\\
&\qquad + c_1\otimes Q(c_3)\otimes c_2 - Q(c_2)\otimes c_1\otimes c_3 + Q(c_3)\otimes c_1\otimes c_2\\
&\quad = Q(Q(c_1)_1)\otimes Q(c_1)_2\otimes c_2 - Q(Q(c_1)_2)\otimes Q(c_1)_1\otimes c_2 - Q(c_1)_1\otimes Q(c_1)_2\otimes c_2\\
&\qquad - Q(Q(c_2)_1)\otimes Q(c_2)_2\otimes c_1 + Q(Q(c_2)_2)\otimes Q(c_2)_1\otimes c_1 + Q(c_2)_1\otimes Q(c_2)_2\otimes c_1\\
&\qquad - Q(c_1)\otimes Q(c_2)\otimes c_3 - Q(c_3)\otimes Q(c_2)\otimes c_1 - Q(c_1)_2\otimes Q(Q(c_1)_1)\otimes c_2\\
&\qquad + Q(c_1)_1\otimes Q(Q(c_1)_2)\otimes c_2 + Q(c_1)_2\otimes Q(c_1)_1\otimes c_2 + Q(c_2)_2\otimes Q(Q(c_2)_1)\otimes c_1\\
&\qquad - Q(c_2)_1\otimes Q(Q(c_2)_2)\otimes c_1 - Q(c_2)_2\otimes Q(c_2)_1\otimes c_1 + Q(c_2)\otimes Q(c_1)\otimes c_3\\
&\qquad + Q(c_2)\otimes Q(c_3)\otimes c_1\\
&\quad \overset{(5)}{=} Q(c_{11})\otimes Q(c_{12})\otimes c_2 - Q(c_{12})\otimes Q(c_{11})\otimes c_2 - Q(c_{21})\otimes Q(c_{22})\otimes c_1\\
&\qquad + Q(c_{22})\otimes Q(c_{21})\otimes c_1 - Q(c_1)\otimes Q(c_2)\otimes c_3 - Q(c_3)\otimes Q(c_2)\otimes c_1\\
&\qquad + Q(c_2)\otimes Q(c_1)\otimes c_3 + Q(c_2)\otimes Q(c_3)\otimes c_1\\
&\quad = 0,
\end{aligned}
$$

finishing the proof. □

Theorem 4.3 Let (C, Δ, R) be a Rota-Baxter coalgebra of weight 0. Define the operation $\widetilde{\Delta}$ on C by

$$\widetilde{\Delta}(c) = Q(c_1)\otimes c_2 - Q(c_2)\otimes c_1.$$

Then $\widetilde{C} = (C, \widetilde{\Delta})$ is a pre-Lie coalgebra.

Proof. Same to the proof of Theorem 4.2. □

5 Rota-Baxter bialgebras

In this section, we will combine the Rota-Baxter algebra and Rota-Baxter coalgebra to Rota-Baxter bialgebra and also provide some examples.

Definition 5.1 Let λ, γ be elements in K and H a bialgebra (maybe without unit and counit). A triple (H, P, Q) is called a Rota-Baxter bialgebra of weight (λ, γ) if (H, P) is a Rota-Baxter algebra of weight λ and (H, Q) is a Rota-Baxter coalgebra of weight γ, that is, H satisfies the following conditions: for all $x, y \in H$,

$$P(x)P(y) = P(xP(y)) + P(P(x)y) + \lambda P(xy),$$

$$(Q \otimes Q)\Delta = (id \otimes Q)\Delta Q + (Q \otimes id)\Delta Q + \gamma \Delta Q.$$

Example 5.2 Let H be a Hopf algebra and (A, μ, Δ) a bialgebra. Suppose that (A, μ) is a right H-Hopf module algebra and (A, Δ) is a right H-Hopf module coalgebra. Then (A, P_R, P_R) is a Rota-Baxter bialgebra of weight $(-1, -1)$.

Proof. By using Theorem 3.2 and [16, Theorem 2.4], we can get the conclusion directly. □

Remark Similarly, the above result holds for the left H-Hopf module algebra and the left H-Hopf module coalgebra.

Example 5.3 Let C be a bialgebra and H a Hopf algebra with the antipode S. Suppose that there are two bialgebra maps: $i : H \longrightarrow C$ and $\pi : C \longrightarrow H$ such that $\pi \circ i = id_H$, i.e., C is a bialgebra with a projection (see [20]). Set $\Pi = id_C \star (i \circ S \circ \pi)$, where $\star$ is the convolution product on $\mathrm{End}(C)$. The right H-Hopf module structure are given by the following:

$$c \cdot h = ci(h),$$

$$\rho_R(c) = c_1 \otimes \pi(c_2),$$

for all $c \in C$ and $h \in H$. Then (C, Π, Π) is a Rota-Baxter bialgebra of weight $(-1, -1)$.

Proof. It is straightforward by using Example 3.3 and [16, Example 2.6]. □

Example 5.4 Let $\{x, y, z\}$ be a basis of a 3-dimensional vector space H over K. We define multiplication and comultiplication over H:

$$x^2 = x, \quad y^2 = y, \quad z^2 = 0$$

$$xy = yx = y, \quad yz = z,$$

$$xz = zx = z, \quad zy = 0,$$

$$\Delta(x) = x \otimes x, \quad \Delta(y) = y \otimes y, \quad \Delta(z) = z \otimes z.$$

It is not hard to see that H is a bialgebra with the above constructions. Let P_1, P_2, Q be operators defined with respect to the basis $\{x, y, z\}$ by

$$P_1(x) = az, \quad P_2(x) = -cx - cy, \quad Q(x) = dz,$$

$$P_1(y) = bz, \quad P_2(y) = -cy, \quad Q(y) = dz,$$

$$P_1(z) = 0, \quad P_2(z) = -cz, \quad Q(z) = 0,$$

where $a, b, c, d \in K$. Then (H, P_1, Q) is a Rota-Baxter bialgebra of weight $(0, d)$, and (H, P_2, Q) is a Rota-Baxter bialgebra of weight (c, d).

Acknowledgments The authors are deeply indebted to the referee for his/her very useful suggestions and some improvements to the original manuscript. This work was partially supported by the Foundation for Key Teacher by the Henan Normal University and the IRTSTHN (No. 14IRTSTHN023).

References

[1] An H H, Bai C M. From Rota-Baxter algebras to pre-Lie algebras. J Phys A, 2008, 41 (1), 015201, 19 pp.

[2] Bai R P, Guo L, Li J Q, Wu Y. Rota-Baxter 3-Lie algebras. J Math Phys, 2013, 54(6), 063504, 14 pp.

[3] Bai C M, Guo L, Ni X. Relative Rota-Baxter operators and tridendriform algebras. J Algebra Appl, 2013, 12(7), 1350027, 18 pp.

[4] Baxter G. An analytic problem whose solution follows from a simple algebraic identity. Pacific J Math, 1960, 10: 731-742.

[5] Bokut L A, Chen Y Q, Qiu J J. Gröbner-Shirshov bases for associative algebras with multiple operators and free Rota-Baxter algebras. J Pure Appl Algebra, 2010, 214 (1): 89-100.

[6] Brzeziński T. Rota-Baxter systems, dendriform algebras and covariant bialgebras, arXiv:1503.05073v1.

[7] Chen Y Q, Mo Q H. Embedding dendriform algebra into its universal enveloping Rota-Baxter algebra. Proc. Amer. Math. Soc. 2011, 139(12): 4207-4216.

[8] Ebrahimi-Fard K, Guo L. Multiple zeta values and Rota-Baxter algebras. Integers, 2008, 8 (2), A4, 18 pp.

[9] Ebrahimi-Fard K, Guo L. Free Rota-Baxter algebras and rooted trees. J Algebra Appl, 2008 7 (2): 167-194.

[10] Ebrahimi-Fard K, Guo L. Rota-Baxter algebras in renormalization of perturbative quantum field theory. Universality and renormalization, 47-105, Fields Inst. Commun., 50, Amer. Math. Soc., Providence, RI, 2007.

[11] Gubarev V Y, Kolesnikov P S. Embedding of dendriform algebras into Rota-Baxter algebras. Cent Eur J Math, 2013, 11 (2): 226-245.

[12] Guo L. An introduction to Rota-Baxter algebra. Surveys of Modern Mathematics, 4. International Press, Somerville, MA; Higher Education Press, Beijing, 2012. xii+226 pp.

[13] Guo L. Properties of free Baxter algebras. Adv Math, 2000, 151: 346-374.

[14] Guo L, Keigher W. Baxter algebras and shuffle products. Adv Math, 2000, 150: 117-149.

[15] Guo L, Zhang B. Polylogarithms and multiple zeta values from free Rota-Baxter algebras. Sci China Math, 2010, 53 (9): 2239-2258.

[16] Jian R Q. Construction of Rota-Baxter algebras via Hopf module algebras. Sci China Math, 2014, 57(11): 2321-2328.

[17] Jian R Q, Zhang J. Rota-Baxter coalgebras, arXiv:1409.3052.

[18] Li X X, Hou D P, Bai C M. Rota-Baxter operators on pre-Lie algebras. J Nonlinear Math Phys, 2007, 14 (2): 269-289.

[19] Montgomery S. Hopf algebras and their actions on rings. CBMS Lectures in Math. Vol. 82, Providence, RI: AMS, 1993.

[20] Radford D E. The structure of Hopf algebra with a projection. J Algebra, 1985, 92: 322-347.

[21] Rota G C. Baxter algebras and combinatorial identities. I, II, Bull Amer Math Soc, 1969, 75: 325-329, 330-334.